

This journal article contains language that may not fall within the scope of FDA clearance of Lipogems and was provided by Lipogems to the requestor in response to an unsolicited request for information. Some authors may have financial interest or receive compensation from the manufacturer. The Lipogems System is a sterile medical device intended for the closed-loop processing of lipoaspirate tissue in medical procedures involving the harvesting, concentrating and transferring of autologous adipose tissue harvested with a legally marketed lipoplasty system. The device is intended for use in the following surgical specialties when the transfer of harvested adipose tissue is desired: orthopedic surgery, arthroscopic surgery, and other specified surgical disciplines.

Biological Stability of Autologous Adipose Tissue Transplantation for Treatment of Hip Chondral Lesions

PURPOSE

The purpose of this study was to evaluate the biochemical factors involved in the chondrogenetic process related to autologous adipose tissue transplantation, for the treatment of chondral lesions in the hip. We analysed the regenerative capacities of the transplant and its effects on a stable biological activity. Our intent was to demonstrate that the autologous adipose tissue transplantation has an intrinsic biological stability and that a mechanical stability of the transplant is not needed to guarantee an effective treatment.

MATERIALS and METHODS

Autologous adipose tissue transplant is obtained from a standard lipoaspirate, processed by clearing both the oily mass and blood residuals and isolating stromal vascular fractions (lipogems®). A mechanical test was performed to compare the compressive strength and the Young elastic modulus of adipose tissue and the transplant. We evaluated the percentage of mesenchymal stem cells present in the transplant, the cellular capacity to produce collagen and GAG and the potentials to chondrogenic differentiation, under the signalling action of TGF- β and FGF-2 factors, as well as the gene expression of protein SOX-9. We also tested the cellular affinity of the transplant to healthy and damaged cartilage and its chondrogenic potentials. Finally we investigated the paracrine effect of autologous adipose tissue transplant on chondrocyte proliferation and cartilage matrix protection using MMP-3 and MMP-13 proteases expression.

RESULTS

Autologous adipose tissue transplant (lipogems®) is a gelatinous viscose tissue, very rich in bio active units called stromal vascular fractions. Mechanical tests demonstrated that this tissue has a significant increase of compressive strength and Young modulus compared to adipose tissue (Fig. 1–2).




Fig. 1 compressive strength




Fig. 2 Young elastic modulus

The concentration of mesenchymal stem cells in the transplant was 2%, therefore much higher compared to the one estimated from bone marrow (0.02%) or from microfractures (0.0002%). Cellular capacity to produce collagen type I/II and GAG was 93% and 90% respectively (fig. 3-4).




Fig. 3 collagen expression.




Fig. 4 GAG expression.

Chondrocytes differentiation, estimated by SOX-9 protein gene expression, was 0.0004 (2-DCT), demonstrating a significant differentiation rate of mesenchymal stem cells (fig. 5).




Fig. 5 Active Chondrocytes marked with SOX-9 Ab.

Cytokines TGF- β and FGF-2 positively influence the chondrogenic differentiation. The transplant demonstrated to have a low or absent cellular affinity and chondrocyte differentiation capacity to normal cartilage (Fig. 6).




Fig. 6 Very few chondrocytes active on a normal chondral surface.




Fig. 7 Increased chondrocytes activity on a damaged chondral surface.

On the contrary cellular affinity and chondrocyte differentiation capacity were very highly expressed on chondral lesions (Fig. 7). MMP-13 proteases expression was significantly lower when associated to autologous adipose tissue transplant, demonstrating a significant paracrine effect.

DISCUSSION

Chondral defects in the hip are very frequent and actually represent a field of major concern for treatment, mainly because biological processes are involved. Biological stability is the expression of all factors involved in a biochemical process resulting in a stable biological activity. In orthopaedics biological stability is opposed to primary stability, which comes from mechanics. Stromal vascular fraction units contained in autologous adipose tissue transplant act as a scaffold for mesenchymal stem cells, enhancing potentials for chondrogenic differentiation. This process appears to be independent from a stable implant and much more correlate to biochemical signals.

CONCLUSIONS

Autologous adipose tissue transplant demonstrated to have a major biological activity and stability to guarantee a regenerative chondrogenetic process in the treatment of hip chondral lesions.

BIBLIOGRAFIA

1. Hunziker E. Articular cartilage repair: basic science and clinical progress. A review of the current status and prospects. *Osteoarthritis Cartilage* 2001; 10: 432-63.
2. Samuels J, Kransnokutsky S. Osteoarthritis: a tale of three tissues. *Bull NYU Hosp* 2008; 66(3):244-50.
3. Sharkey PF, Cohen SB, Leinberry CF, Parvizi J. Subchondral bone marrow lesions associated with knee osteoarthritis. *Am J Orthop* 2012; 41(9):413-7.
4. Breyner N, Hell R, Carvalho L, Machado C, Peixoto IN, Valerio P, et al. Effect of a three-dimensional Chitosan porous scaffold on the differentiation of mesenchymal stem cells into Chondrocytes. *Cells Tissues Organs* 2010; 191(2):119-28.
5. Fukuda K. Intra-articular injection of hyalur for treatment of knee osteoarthritis. *Clin Calcium* 2004; 14(7):103-7.
6. Bendinelli P, et al. Molecular basis of anti-inflammatory action of platelet-rich plasma on human chondrocytes: mechanisms of NF-kB inhibition via HGF. *J Cell Physiol* 2010; 225(3):757-66.
7. Leunig M, Tibor LM, Naal FD, Ganz R, Steinwachs MR. Surgical technique: second generation bone marrow stimulation via surgical dislocation to treat hip cartilage lesions. *Clin Orthop Relat Res* 2012; 470(12):3421-31.
8. Kuroda T, Matsumoto T, Mifune Y, Fukui T, Kubo S, Matsushita T, et al. Therapeutic strategy of third-generation autologous chondrocyte implantation for osteoarthritis. *Ups J Med Sci* 2011;116(2):107-14.
9. Peterson L, Minas T, Brittberg M, Lindahl A. Treatment of osteochondritis dissecans of the knee with autologous chondrocyte transplantation: results at two to ten years. *J Bone Joint Surg Am* 2003; 85A(Suppl 2): 17-24.
10. Emans PJ, Jansen EJ, van Iersel D, Welting TJ, Woodfield TB, Bulstra SK, et al. Tissue-engineered constructs: the effect of scaffold architecture in osteochondral repair: *J Tissue Eng Regen Med* 2013; 7(9):751-6.
11. Sakaguchi Y, Sekiya I, Yagishita K, Muneta T. Comparison of human stem cells derived from various mesenchymal tissues: superiority of synovium as a cell source. *Arthritis Rheum* 2005; 52:2521-9.
12. Xue JX, Gong YY, Zhou GD, Liu W, Cao Y, Zhang WJ. Chondrogenic differentiation of bone marrow-derived mesenchymal stem cells induced by acellular cartilage sheets. *Biomaterials* 2012; 33:5832-40.
13. Freyria AM, Mallein-Gerin F. Chondrocytes or adult stem cells for cartilage repair: the indisputable role of growth factors. *Injury, Int J Care Injured* 2012; 43:259-65.
14. Kastrinaki MC, Andreakou I, Charbord P, Papadaki HA. Isolation of human bone marrow mesenchymal stem cells using different membrane markers: comparison of colony/cloning efficiency, differentiation potential, and molecular profile. *Tissue Eng Part C Methods* 2008; 14:333-9.

15. Shenaq D, Rastegar F, Petkovic D, Zhang BQ, He BC, Chen L, et al. Mesenchymal progenitor cells and their orthopaedic applications: forging a path towards clinical trials. *Stem Cells Int* 2010; 519028
16. www.tursiops-biology.com/gfp.htm
17. M. Ian Phillips. *Methods in enzymology*. 346(26): 454-465.(2002)
18. Ikeda T., Kamekura S., Mabuchi A., Kou I., Seki S., Takato T., Nakamura K., Kawaguchi H., Ikegawa S., Chung U., The Combination of SOX5, SOX6, and SOX9 (the SOX Trio) Provides Signals Sufficient for Induction of Permanent Cartilage, *Arthritis & Rheumatism*: 50 (11): 3561-3573 (2004)
19. Grigolo B., Roseti L., Fiorini M., De Franceschi L., Facchini A., Applicazioni dell'ingegneria tissutale: riparazione di lesioni cartilaginee con condrociti autologhi, *Reumatismo*: 54 (4): 364-371 (2002)
20. www.leicabiosystems.com
21. Ramirez G.A., Altimira J., Garcia B., Vilafranca M., Chondro-osteoblastic Metaplasia in Canine Benign Cutaneous Lipomas, *Journal of Comparative Pathology*: 142: 89-93 (2010)
22. Van Lent P., Van den Bergmesenchimal W., Stem cell therapy in osteoarthritis: advanced tissue or intervention with smouldering synovial activation?, *Arthritis research&therapy*: 15 (112).