



Treatment of Achilles Tendon Partial Tear with Lipoaspirate (Lipogems®)

Brady Saray-Logan, Christopher J Rogers MD, RMSK
San Diego Orthobiologics Medical Group, Carlsbad, CA

ABSTRACT

Elite and recreational athletes commonly present with Achilles tendinopathy and partial tears. The tendon pathology demonstrates a failed healing response rather than an inflammatory process. The source of pain in tendinopathy is typically related to the neurovascular ingrowth seen in the tendon's response to injury^{3,4}. Conservative medical treatments include rest, bracing, eccentric strength training, low energy shock wave therapy, topical glyceryl trinitrate⁶. More recently, minimally invasive ultrasound guided treatments such as tendon fenestration, tendon scraping and hydro-dissection of the paratenon have been reviewed. Surgical intervention is rarely indicated.

Recent studies have demonstrated that lipoaspirate contains and produces growth-factors, such as platelet-derived growth-factor (PDGF), fibroblast growth-factor (FGF), transforming growth-factor beta (TGF- β), and vascular endothelial growth-factor (VEGF)¹. These growth factors are known to play an important regulatory role in cellular proliferation, cellular migration, matrix synthesis and angiogenesis. The Lipogems® system can produce a non-expanded, rinsed, and microfragmented adipose tissue graft that maintains an intact stromal vascular niche and contains cellular elements with mesenchymal stem cell and pericyte characteristics¹¹.

We present the case of a 16 year old girl who developed Achilles tendinopathy and partial tear that was non-responsive to conservative care. She was treated with ultrasound guided lipoaspirate (Lipogems®) injection. We report her significant improvement in pain and physical function at six month follow-up. Future clinical trials of lipoaspirate for the treatment of chronic Achilles tendinopathy and tear are warranted.

INTRODUCTION

Chronic degenerative tendon disorders (tendinopathy) occur frequently and are difficult to treat. Tendon injuries are often associated with significant physical dysfunction and disability, due to the limited self-repair capacity and propensity for scar formation¹. Non-insertional Achilles tendinopathy is a common cause of ankle pain and typically occurs 2 - 6 cm proximal to the tendon insertion on the calcaneus.

Athletes, whether elite or recreational, are the most common group to present with Achilles tendinopathy, but it is also found in people with advanced age, obesity, diabetes or hypertension. Other risk factors for Achilles tendinopathy include previous tendon injury, decreased muscle strength, altered gait kinematics, limited ankle dorsiflexion, training errors, and the use of steroids or fluoroquinolones.

The degenerative process is essentially a failed healing response in the tendon². Fibromatous degeneration is the most frequent finding and is related to the relative hypovascularity of the critical zone. Mucoid degeneration is the second most common type of degeneration with large mucoid patches and vacuoles found between thinned degenerated tendon fibers. Early episodes are asymptomatic, however symptoms develop when the vacuoles and lacunae coalesce to form an interstitial tear. Histological degenerative changes include loss of the normal collagenous architecture, replacement with amorphous mucinous material, hypercellularity, increased glycosaminoglycan content and neovascularization. Less commonly, tenolipomatosis or calcifying tendinopathy with tendon calcification may occur³.

Patients often present with focal swelling and tenderness to palpation on the posteromedial aspect of the tendon. Both ultrasound and magnetic resonance imaging (MRI) assist in the diagnosis of Achilles tendinopathy, but ultrasound has a higher degree of accuracy and is more cost effective⁴. Ultrasound imaging may reveal a fusiform tendon with peritendineum thickening, hypoechoic foci, intrasubstance tendon tears, decreased tendon gliding and intratendinous and peritendinous vascularity on color Doppler⁵.

Conservative care of Achilles tendinopathy includes removal of precipitating factors, rest, training routine modifications and the use of orthotics to correct foot or ankle malalignment. Decreased myotendinous flexibility may be treated with stretching. Muscle weakness is treated with progressive eccentric strengthening exercises twice a day for up to 12 weeks using either the Alfredson or Silbernagel protocol. Combining these protocols with low-energy shock-wave therapy (ESWT) may improve outcomes⁶.

The combination of autologous platelets and a fibrin matrix in a sheep tendinopathy model has demonstrated increased tenocyte proliferation, synthesis of type-I collagen and angiogenic factors (VEGF, HGF) promoting neovascularization⁷. In a prospective clinical case series of 14 patients with chronic non-insertional Achilles tendinopathy, leukocyte-rich platelet rich plasma (LR-PRP) demonstrated significant improvement in the American Orthopedic Foot and Ankle Society (AOFAS) scale, the Victorian Institute of Sport Assessment – Achilles (VISA-A) scale and ultrasound imaging results at 18 month followup⁸. A well-designed RCT study of chronic Achilles tendinopathy found no significant difference in pain or activity level 24 weeks after receiving an ultrasound guided LR-PRP injection combined with an eccentric stretching program⁹. Yet, in a systematic review and meta-analysis, LR-PRP demonstrated a strongly positive effect when compared with LP-PRP, suggesting that what is injected and how it is injected may influence outcomes⁹.

Up to 33% of patients will fail conservative care and consider surgical excision of the pathologic tissue with or without tendon transfer augmentation. Open surgery is associated with an overall complication rate of 11% and success rate ranging from 50-100%. Minimally-invasive percutaneous therapies designed to strip the paratenon from the tendon with high-volume fluid injection or ventral tendon scraping have shown promise in relieving symptoms. Percutaneous tendon fenestration (tenotomy) performed under ultrasound guidance has also demonstrated good outcomes.

Adipose tissue is a connective tissue derived from embryonic mesoderm, consisting of a heterogeneous population of cells such as adipocytes, preadipocytes, smooth muscle cells, endothelial cells, mast cells and fibroblast immune cells. From adipose tissue manipulation it is possible to isolate these so called stromal vascular fraction (SVF) which contains, among others, mesenchymal stem cells (MSC)¹⁰. When compared with bone marrow, adipose tissue contains several times the number of pluripotent stem cells^{12,13}.

CASE REVIEW

Our patient was a 16 year old female student athlete who presented with a two year history of progressive right heel pain. She notes history of several injuries to the ankle while playing basketball. One particular day, she noticed a "pop" associated with increased Achilles tendon pain. She had pain with many activities, but mostly when walking around campus or climbing stairs. She was a basketball player, but had been unable to practice or exercise for more than one year. She was using a CAM walker boot for almost a year while walking or performing activities of daily living. She had been diagnosed with chronic non-insertional Achilles tendinopathy on the basis of MRI and clinical presentation. She had received more than 7 months of physical therapy which included eccentric calf strengthening, soft tissue mobilization, and modalities. She did not use oral analgesics, NSAIDs, or topical medications. She had not received any form of injection therapy. She was using acetaminophen 500mg and tramadol 50mg for pain intermittently. She did not take any other medications. She denied the use of antibiotics in the past. She denied any allergies or other medical conditions. She has a family history of arthritis, but denied any family history of autoimmune disease, connective tissue disease or seronegative arthritis.


On physical examination she presented wearing her CAM walker boot. She had significant focal swelling of the right Achilles tendon which was tender to light palpation. The calf squeeze test was negative. A 4 cm palpable, firm and tender mass was located from 2 cm proximal to the tendon insertion. Ankle range of motion was limited to 20 degrees dorsiflexion and 20 degrees plantarflexion. She was able to actively move the ankle, but was unable to hop, squat or perform a single heel raise. Her light touch sensation was intact and the dorsalis pedis and posterior tibial pulses were intact in the lower extremity. She was noted to have moderate pes planus bilaterally, but normal subtalar joint motion, leg length and triceps surae muscle bulk.

Her Foot and Ankle Disability Index Score (FADI) was 65.4 and her average Visual Analog Score (VAS) was 7 out of 10.

Magnetic resonance imaging (MRI) of the right ankle revealed Achilles tendinosis with a 3.3 cm longitudinal split tear, extending from 2.7 cm to 6 cm proximal to the tendon insertion.

Musculoskeletal Ultrasound (MSK US) Examination of the right ankle was performed using a 8-12 MHz linear probe. Examination revealed fusiform thickening, diffuse hypoechoic defects and hypoechoic disorganization of the tendon fibrillar pattern from 2 to 6 cm proximal to the Achilles tendon insertion on the calcaneus. Edema was evident in Kager's fat pad anterior to the Achilles tendon. Mobility of the Achilles tendon with ankle motion was limited. The retrocalcaneal bursa, plantaris tendon and FHL tendon were unremarkable.

Given the radiographic findings, duration of symptoms and level of functional limitations, we elected to treat with lipoaspirate injection (Lipogems®) of the Achilles tendon. The Lipogems® system can produce a non-expanded, rinsed, and microfragmented adipose tissue graft that maintains an intact stromal vascular niche and contains cellular elements with mesenchymal stem cell and pericyte characteristics. We hypothesized that these cells would express a phenotype that could encourage tendon neovascularization, tenocyte proliferation and type-I collagen synthesis. Platelet rich plasma has also demonstrated increased tenocyte proliferation, synthesis of type-I collagen and promotion of neovascularization. However, in this case, we believed that the degree of tissue degeneration and the duration of symptoms warranted the addition of adipose derived cells to augment any benefit provided by platelet rich plasma. Studies to inform the physician as to the number or type of cells required to heal such an injury do not currently exist. We were limited by the volume that can be injected into a degenerated tendon and adjusted our treatment plan accordingly.



Performing lipoaspiration with 13 G cannula and VacLok syringe.



Preparation of the lipoaspirate using the Lipogems® system.

PROCEDURE

The patient was informed of all other possible treatment options and informed consent was obtained from the patient and her mother to proceed with PRP / lipoaspirate injection. The patient was advised to avoid the use of aspirin or NSAIDs for one week prior to the procedure. On the day of the procedure, she was given Ancef 1 gm IVP 60 minutes prior to the procedure. Fifty-two milliliters of blood was obtained from the antecubital fossa to prepare 3 ml of leukocyte poor PRP using the 1% hematocrit setting on the PRP processing centrifuge (Arthrex Angel, Naples FL). We did not determine the baseline serum platelet level or assay the PRP sample to determine platelet content.

The patient was placed in the prone position in our procedure suite and monitored with blood pressure, pulse oximetry and electrocardiographic monitoring. The skin over her lateral flank was prepped with Hibiclen and sterilely draped. The skin was anesthetized with 5 ml 1% lidocaine. A 2 mm stab incision was made with an #11 blade. Tumescant anesthesia was performed with 96 ml Klein's solution and a 17Gx185 mm anesthetic infiltration cannula. Klein's solution had been prepared with 500 ml normal saline, 20 ml of 2% lidocaine and 0.5 ml 1:1000 epinephrine. After waiting 12 minutes for detumescence, lipoaspiration was performed with a 13Gx185 mm cannula and VacLok syringe to produce 20 ml lipoaspirate. At this time, the Lipogems® method was used to process the lipoaspirate. In brief, the Lipogems® method allows for a minimal enzyme-free manipulation in a sealed, sterile device to minimize trauma to the cellular content. The device contained normal saline and filters which gradually reduce the adipose clusters and eliminates oil, hematic residue and tumescent anesthesia. We obtained 8ml of washed and rinsed lipoaspirate (Lipogems®) for injection. The patient's wound was cleansed and dressed.

The ankle was prepped with Hibiclen and sterilely draped. The skin over the medial aspect of the Achilles tendon was anesthetized with 1% lidocaine at two locations. The tendon and paratenon were not anesthetized. The tendon was visualized with ultrasound and a sterilely draped 12 MHz linear probe. An 18 g needle was advanced into the tendon to perform tenotomy at multiple locations with special attention to treat the hypoechoic aspects of the tendon until the consistency of the tendon was softened. Using the 18 g needle and 1ml syringe containing 0.5 ml of Lipogems® was injected at 8 locations throughout the tendon without resistance. Lipogems® was also injected anterior and medial to the tendon. After injection of the Lipogems®, 3 ml of LP-PRP was injected at several locations within the tendon. The injection site was cleansed and bandaged. The patient presented to our recovery room and was discharged within 20 minutes. She was advised to wear her CAM walker boot and to use ice, tylenol or Tramadol as needed for up to three days. She was given an abdominal binder and prescription for cephalexin 250 mg QID for 5 days.

RECOVERY


The patient presented to our clinic for followup one week after her procedure. She denied any complications, side effects or significant pain. Her incision site had produced clear discharge for one day and was healed in three days. She did not have any bruising of the adipose harvest site. Her ankle was stiff and swollen, but not red or significantly tender. She was able to walk short distances with the CAM walker boot. She was instructed to start physical therapy which included passive/active range of motion, edema management, soft tissue mobilization and progression of strength training as tolerated. She was allowed to bear weight as tolerated.

At six weeks followup, our patient noticed a significant reduction in her pain at rest and with ambulating short distances. She was not compliant with physical therapy, choosing to stretch and ambulate on her own schedule. She had developed mild kinesophobia due to the duration of her injury. On physical examination, the tendon swelling has reduced and the tendon was less tender to palpation. Ultrasound examination revealed an improvement in the hyperechoic fibrillar pattern of the tendon, however the tendon thickening and hypoechoic defects remained. The patient was encouraged to initiate the formal physical therapy program.

At three month followup, our patient noticed substantial reduction in pain and improved physical activity. Her repeat FADI score was 97.1. Her average VAS was zero out of ten. She could walk, climb and perform heel raises and squats without pain. On physical examination, she had full dorsiflexion and normal plantar flexion strength. The tendon was not tender to palpation. Ultrasound examination revealed significant improvement in the hyperechoic fibrillar pattern of the tendon with near complete resolution of the hypoechoic defects. She was compliant with an eccentric strengthening program in physical therapy twice a week and at home twice a week.

At six month followup, our patient is able to walk long distances and climb stairs without pain. She can run up to 30 minutes at a time and performs all strengthening exercises without pain. Although not required, we elected to repeat her ankle MRI which revealed resolution of the high intensity linear band and diffuse speckled signal consistent with resolution of her tendinopathy and partial tears. The tendon thickness and Kager's fat pad edema was improved as well.


At this time, she takes no medication and does not wear a CAM walker boot. But she does have plans to attend her high school prom and wear high heels.



Sagittal ultrasound imaging reveals fusiform thickening, hypoechoic defects, and disorganization of fibrillar pattern in the Achilles tendon.




Sagittal (STIR) MRI image shows 3.3 cm longitudinal partial tear in Achilles tendon.



Sagittal ultrasound imaging reveals improved fusiform thickening, resolution of hypoechoic defects and increased hyperechoic fibrillar pattern of the Achilles tendon 6 months after treatment



Sagittal (STIR) MRI image reveals resolution of partial tear of the Achilles tendon 6 months after treatment



Mesenchymal stem cell (MSC) emerging from lipoaspirate

Foot & Ankle Disability Index (FADI)	PRE score: 65.4	3 month POST score: 97.1
1. Pain		10
2. Stair		10
3. Walk		10
4. Run		10
5. Climb		10
6. Jump		10
7. Sit		10
8. Stand		10
9. Bend		10
10. Push		10
11. Pull		10
12. Twist		10
13. Lift		10
14. Carry		10
15. Throw		10
16. Kick		10
17. Bend		10
18. Stand		10
19. Sit		10
20. Walk		10
21. Run		10
22. Climb		10
23. Jump		10
24. Bend		10
25. Stand		10
26. Sit		10
27. Walk		10
28. Run		10
29. Climb		10
30. Jump		10
31. Bend		10
32. Stand		10
33. Sit		10
34. Walk		10
35. Run		10
36. Climb		10
37. Jump		10
38. Bend		10
39. Stand		10
40. Sit		10
41. Walk		10
42. Run		10
43. Climb		10
44. Jump		10
45. Bend		10
46. Stand		10
47. Sit		10
48. Walk		10
49. Run		10
50. Climb		10
51. Jump		10
52. Bend		10
53. Stand		10
54. Sit		10
55. Walk		10
56. Run		10
57. Climb		10
58. Jump		10
59. Bend		10
60. Stand		10
61. Sit		10
62. Walk		10
63. Run		10
64. Climb		10
65. Jump		10
66. Bend		10
67. Stand		10
68. Sit		10
69. Walk		10
70. Run		10
71. Climb		10
72. Jump		10
73. Bend		10
74. Stand		10
75. Sit		10
76. Walk		10
77. Run		10
78. Climb		10
79. Jump		10
80. Bend		10
81. Stand		10
82. Sit		10
83. Walk		10
84. Run		10
85. Climb		10
86. Jump		10
87. Bend		10
88. Stand		10
89. Sit		10
90. Walk		10
91. Run		10
92. Climb		10
93. Jump		10
94. Bend		10
95. Stand		10
96. Sit		10
97. Walk		10
98. Run		10
99. Climb		10
100. Jump		10
101. Bend		10
102. Stand		10
103. Sit		10
104. Walk		10
105. Run		10
106. Climb		10
107. Jump		10
108. Bend		10
109. Stand		10
110. Sit		10
111. Walk		10
112. Run		10
113. Climb		10
114. Jump		10
115. Bend		10
116. Stand		10
117. Sit		10
118. Walk		10
119. Run		10
120. Climb		10
121. Jump		10
122. Bend		10
123. Stand		10
124. Sit		10
125. Walk		10
126. Run		10
127. Climb		10
128. Jump		10
129. Bend		10
130. Stand		10
131. Sit		10
132. Walk		10
133. Run		10
134. Climb		10
135. Jump		10
136. Bend		10
137. Stand		10
138. Sit		10
139. Walk		10
140. Run		10
141. Climb		10
142. Jump		10
143. Bend		10
144. Stand		10
145. Sit		10
146. Walk		10
147. Run		10
148. Climb		10
149. Jump		10
150. Bend		10
151. Stand		10
152. Sit		10
153. Walk		10
154. Run		10
155. Climb		10
156. Jump		10
157. Bend		10
158. Stand		10
159. Sit		10
160. Walk		10
161. Run		10
162. Climb		10
163. Jump		10
164. Bend		10
165. Stand		10
166. Sit		10
167. Walk		10
168. Run		10
169. Climb		10
170. Jump		10
171. Bend		10
172. Stand		10
173. Sit		10
174. Walk		10
175. Run		10
176. Climb		10
177. Jump		10
178. Bend		10
179. Stand		10
180. Sit		10
181. Walk		10
182. Run		10
183. Climb		10
184. Jump		10
185. Bend		10
186. Stand		10
187. Sit		10
188. Walk		10
189. Run		10
190. Climb		10
191. Jump		10
192. Bend		10
193. Stand		10
194. Sit		10
195. Walk		10
196. Run		10
197. Climb		10
198. Jump		10
199. Bend		10
200. Stand		10
201. Sit		10
202. Walk		10
203. Run		10
204. Climb		10
205. Jump		10
206. Bend		10
207. Stand		10
208. Sit		10
209. Walk		10
210. Run		10
211. Climb		10
212. Jump		10
213. Bend		10
214. Stand		10
215. Sit		10
216. Walk		10
217. Run		10
218. Climb		10
219. Jump		10
220. Bend		10
221. Stand		10
222. Sit		10
223. Walk		10
224. Run		10
225. Climb		10
226. Jump		10
227. Bend		10
228. Stand		10
229. Sit		10
230. Walk		10
231. Run		10
232. Climb		10
233. Jump		10
234. Bend		10
235. Stand		10
236. Sit		10
237. Walk		10
238. Run		10
239. Climb		10
240. Jump		10
241. Bend		10
242. Stand		10
243. Sit		10
244. Walk		10
245. Run		10
246. Climb		10
247. Jump		10
248. Bend		10
249. Stand		10
250. Sit		10
251. Walk		10
252. Run		10
253. Climb		10
254. Jump		10
255. Bend		10
256. Stand		10
257. Sit		10
258. Walk		10
259. Run		10
260. Climb		10
261. Jump		10
262. Bend		10
263. Stand		10
264. Sit		10
265. Walk		10
266. Run		10
267. Climb		10
268. Jump		10
269. Bend		10
270. Stand		10
271. Sit		10
272. Walk		10
273. Run		10
274. Climb		10
275. Jump		10
276. Bend		10
277. Stand		10
278. Sit		10
279. Walk		10
280. Run		10
281. Climb		10
282. Jump		10
283. Bend		10
284. Stand		10
285. Sit		10
286. Walk		10
287. Run		10
288. Climb		10
289. Jump		10
290. Bend		10
291. Stand		10
292. Sit		10
293. Walk		10
294. Run		10
295. Climb		10
296. Jump		10
297. Bend		10
298. Stand		10
299. Sit		10
300. Walk		10
301. Run		10
302. Climb		10
303. Jump		10
304. Bend		10
305. Stand		10
306. Sit		10
307. Walk		10
308. Run		10
309. Climb		10
310. Jump		10
311. Bend		10
312. Stand		10
313. Sit		10
314. Walk		10
315. Run		10
316. Climb		10
317. Jump		10
318. Bend		10
319. Stand		10
320. Sit		10
321. Walk		10
322. Run		10
323. Climb		10
324. Jump		10
325. Bend		10
326. Stand		10
327. Sit		10
328. Walk		10
329. Run		10
330. Climb		10
331. Jump		10
332. Bend		10
333. Stand		10
334. Sit		10
335. Walk		10
336. Run		10
337. Climb		10
338. Jump		10
339. Bend		10
340. Stand		10
341. Sit		10
342. Walk		10
343. Run		10
344. Climb		10
345. Jump		10
346. Bend		10
347. Stand		10
348. Sit		10
349. Walk		10
350. Run		10
351. Climb		10
352. Jump		10
353. Bend		10
354. Stand		10
355. Sit		10
356. Walk		10
357. Run		10
358. Climb		10
359. Jump		10
360. Bend		10
361. Stand		10
362. Sit		10
363. Walk		10
364. Run		10
365. Climb		10
366. Jump		10
367. Bend		10
368. Stand		10
369. Sit		10
370. Walk		10
371. Run		10
372. Climb		10
373. Jump		10
374. Bend		10
375. Stand		10
376. Sit		10
377. Walk		10
378. Run		10
379. Climb		10
380. Jump		10
381. Bend		10
382. Stand		10
383. Sit		10
384. Walk		10
385. Run		10
386. Climb		10
387. Jump		10
388. Bend		10
389. Stand		10
390. Sit		10
391. Walk		10
392. Run		10
393. Climb		10
394. Jump		10
395. Bend		10
396. Stand		10
397. Sit		10
398. Walk		10
399. Run		10
400. Climb		10
401. Jump		10
402. Bend		10
403. Stand		10
404. Sit		10
405. Walk		10
406. Run		10
407. Climb		10
408. Jump		10
409. Bend		10
410. Stand		10
411. Sit		10
412. Walk		10
413. Run		10
414. Climb		10
415. Jump		10
416. Bend		10
417. Stand		10
418. Sit		10
419. Walk		10
420. Run		10
421. Climb		10
422. Jump		10
423. Bend		10
424. Stand		10
425. Sit		10
426. Walk		10
427. Run		10
428. Climb		10
429. Jump		10
430. Bend		10
431. Stand		10
432. Sit		10
433. Walk		10
434. Run		10
435. Climb		10
436. Jump		10
437. Bend		10
438. Stand		10
439. Sit		10
440. Walk		10
441. Run		10
442. Climb		10
443. Jump		10
444. Bend		10
445. Stand		10
446. Sit		10
447. Walk		10
448. Run		10
449. Climb		10
450. Jump		10
451. Bend		10
452. Stand		10
453. Sit		10
454. Walk		10
455. Run		10
456. Climb		10
457. Jump		10
458. Bend		10
459. Stand		10
460. Sit		10
461. Walk		10
462. Run		10
463. Climb		10
464. Jump		10
465. Bend		10
466. Stand		10
467. Sit		10
468. Walk		10
469. Run		10
470. Climb		10
471. Jump		10
472. Bend		10
473. Stand		10
474. Sit		10
475. Walk		10
476. Run		10
477. Climb		10
478. Jump		10
479. Bend		10
480. Stand		10
481. Sit		10
482. Walk		10
483. Run		10
484. Climb		10
485. Jump		10
486. Bend		10
487. Stand		10
488. Sit		10
489. Walk		10
490. Run		10
491. Climb		10
492. Jump		10
493. Bend		10
494. Stand		10
495. Sit		10
496. Walk		10
497. Run		10
498. Climb		10
499. Jump		10
500. Bend		10
501. Stand		10
502. Sit		10
503. Walk		10
504. Run		10
505. Climb		10
506. Jump		10
507. Bend		10
508. Stand		10
509. Sit		10
510. Walk		10
511. Run		10
512. Climb		10
513. Jump		10
514. Bend		10
515. Stand		10